

## TRACE ELEMENTS AND OXY-PAH IN SOUTH AMERICAN URBAN SITES: SÃO PAULO, LIMA E MEDELLIN

Guilherme M. Pereira; Huang Xian, Rajasekhar Balasubramanian, Odon Roman Sánchez-Ccoyllo, Nestor Rojas, Pérola de C. Vasconcellos

*University of São Paulo, Institute of Chemistry, Av. Prof. Lineu Prestes, 748, 05508-000, São Paulo – SP, Brazil.*

E-mail: [martinspereira2@hotmail.com](mailto:martinspereira2@hotmail.com)

---

**Objectives:** The South American Emissions Megacities and Climate (SAEMC) project focused on the investigation of air composition and its impact on climate over South American cities; among them: São Paulo (SPA, Brazil), Lima (LIM, Peru) and Medellin (MED, Colombia). São Paulo site is the largest South American city, with 19 million inhabitants, Lima is a coastal city that counts with 8 million inhabitants and Medellin is a 2.5 million inhabitants city, located in a valley. **Methodology:** Particulate matter (PM<sub>10</sub>) was collected in quartz fiber filters with a high volume sampler, the intensive sampling was carried out in June, 2010. Oxy-PAHs were determined by gas chromatography and trace elements were determined by inductively coupled plasma mass spectrometry.

**Results:** The quinone fraction of PAHs have been attributed to toxic effects to health; 9,10-anthraquinone (AQ) is classified as possibly carcinogenic to humans (2B)<sup>1,2</sup>. 9,10-Anthraquinone (AQ) and 9-fluorenone (FO) can originate from both primary and secondary sources<sup>3</sup>. AQ had higher concentrations in MED, while FO had higher concentrations in LIM. Benz[a]anthracene-7,12-dione (BaAQ) concentrations were higher in SPA; BaAQ is attributed to the ozonation of precursor PAH<sup>4</sup>. Concentrations of potassium were higher in SPA and MED (805, 873 and 402 ng m<sup>-3</sup>, for SPA, MED and LIM, respectively); suggesting higher biomass burning sources. Zinc, often associated with industrial emissions and traffic (tyre abrasion), was higher in SPA and lower in LIM (350, 205 and 164 ng m<sup>-3</sup>, for SPA, MED and LIM, respectively). Copper (wear emissions) was also higher in SPA (123, 42 and 17 ng m<sup>-3</sup> for SPA, MED and LIM, respectively). Iron was higher in SPA and attributed to soil resuspension and road dust (1761, 632 and 740 ng m<sup>-3</sup> for SPA, MED and LIM, respectively). **Conclusions:** Vehicular and biomass burning sources appeared to be more important for both SPA and MED sites.

1. IARC. List Of Classification. [http://monographs.iarc.fr/ENG/Classification/List\\_of\\_Classifications.pdf](http://monographs.iarc.fr/ENG/Classification/List_of_Classifications.pdf). Published 2017.
2. Kennedy IM. The health effects of combustion-generated aerosols. *Proc Combust Inst.* 2007;31:2757-2770. doi:10.1016/j.proci.2006.08.116.
3. Souza KF, Carvalho LRF, Allen AG, Cardoso AA. Diurnal and nocturnal measurements of PAH, nitro-PAH, and oxy-PAH compounds in atmospheric particulate matter of a sugar cane burning region. *Atmos Environ.* 2014;83:193-201. doi:10.1016/j.atmosenv.2013.11.007.
4. Ringuet J, Albinet A, Leoz-Garziandia E, Budzinski H, Villenave E. Diurnal/nocturnal concentrations and sources of particulate-bound PAHs, OPAHs and NPAHs at traffic and suburban sites in the region of Paris (France). *Sci Total Environ.* 2012;437:297-305. doi:10.1016/j.scitotenv.2012.07.072.